

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/GB05/000459

International filing date: 11 February 2005 (11.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: GB  
Number: 0403041.7  
Filing date: 11 February 2004 (11.02.2004)

Date of receipt at the International Bureau: 16 March 2005 (16.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



PCT/GB2005/000459



INVESTOR IN PEOPLE


The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed


*Stephen Hordley*

Dated 7 March 2005

100

100

100



# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

1. Your reference

AHB/CP6190722

12FEB04 E872603-1 000060  
P01/7700 0.00-0403041.7 NONE

2. Patent application number

(The Patent Office will fill this part in)

0403041.7

11 FEB 2004

3. Full name, address and postcode of the or of each applicant (underline all surnames)

MILNER, ANNE JOSEPHINE  
YCR P53 RESEARCH GROUP  
DEPARTMENT OF BIOLOGY (AREA 14)  
YORK  
YO10 5DD

Patents ADP number (if you know it)

8189383002

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention

INDUCTION OF APOPTOSIS

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

MEWBURN ELLIS  
York House  
23 Kingsway  
London WC2B 6HP

Patents ADP number (if you know it)

109006

6. Priority: Complete this section if you are declaring priority from one or more earlier patent applications, filed in the last 12 months.

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. Divisionals, etc.: Complete this section only if this application is a divisional application or resulted from an entitlement dispute (see note f)

Number of earlier UK application

Date of filing  
(day / month / year)

8. Is a Patents Form 7/77 (Statement of inventorship and of right to grant of a patent) required in support of this request?

YES


Answer YES if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

Otherwise answer NO (See note d)

Patents Form 1/77

9. Accompanying documents: A patent application must include a description of the invention. Not counting duplicates, please enter the number of pages of each item accompanying this form:

Continuation sheets of this form	0
Description	36
Claim(s)	3
Abstract	0
Drawing(s)	27 + 22 

10. If you are also filing any of the following, state how many against each item.

Priority documents	0
Translations of priority documents	0
Statement of inventorship and right to grant of a patent (Patents Form 7/77)	0
Request for a preliminary examination and search (Patents Form 9/77)	0
Request for a substantive examination (Patents Form 10/77)	0
Any other documents (please specify)	0

11. I/We request the grant of a patent on the basis of this application.

Signature(s)



Date 10 February 2004

12. Name, daytime telephone number and e-mail address, if any, of person to contact in the United Kingdom

ADRIAN H BRASNETT

020 7240 4405

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered YES in part 8, a Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- Part 7 should only be completed when a divisional application is being made under section 15(4), or when an application is being made under section 8(3), 12(6) or 37(4) following an entitlement dispute. By completing part 7 you are requesting that this application takes the same filing date as an earlier UK application. If you want the new application to have the same priority date(s) as the earlier UK application, you should also complete part 6 with the priority details.

## Induction of apoptosis

### Field of the Invention

This invention relates to the induction of apoptosis by inhibition of SIRT1 expression, in particular the induction of apoptosis in tumour cells. Materials and methods for inhibiting SIRT1 expression are provided, including RNA interference methods.

### Background to the Invention

Many different types of stress stimulate cellular signalling pathways that result in stabilisation and activation of the tumour suppressor p53 (reviewed in Pluquet and Hainaut, 2001). Stabilisation of p53 is invariably accompanied by extensive post-translational modifications, including phosphorylation and acetylation (see Appella and Anderson, 2001). Mapping the precise relationships between stress stimuli, specific modifications and the stabilisation and activation of p53 has proven extremely difficult; however general patterns and more specific correlations are now established, particularly with regard to phosphorylation of the amino-terminus and acetylation of the carboxy-terminus.

The transcriptional activity of p53, with particular regard to its pro-apoptotic functions, is tightly regulated. Therefore if acetylation functions to activate p53 transcriptional activity, a logical assumption would be that p53 acetylation is subject to negative control.

Until recently, deacetylation of p53 was only known to be performed by members of the trichostatin A-sensitive histone

deacetylase (HDAC) class I family (Juan et al., 2000; Luo et al., 2000). Indeed, mounting evidence suggests that p53 utilises these HDACs to repress specific promoters (Murphy et al., 1999). More recently, the human sirtuin SIRT1 (Frye, 1999) has been identified as a *bona fide* p53 deacetylase (Luo et al., 2001; Vaziri et al., 2001).

The sirtuins are a ubiquitous gene family found throughout eukarya and prokarya, defined by conserved ~250 amino acid core domain. Many of the sirtuins are NAD-dependent deacetylases ['NDAC'].

The function of *Saccharomyces cerevisiae* sirtuin SIR2 has been extensively studied. It has many activities including silencing of mating-type loci, telomeric position effect silencing and silencing at the rDNA locus, suppression of illegitimate recombination and increasing cellular control of longevity. It is also implicated in response to dsDNA breaks.

Humans have seven sirtuins, although not all appear to have NDAC activity.

Human SIRT2 is a cytoplasmic, microtubule-associated protein. It shows increases in abundance and phosphorylation at G2/M. It is a tubulin deacetylase and is strongly down-regulated in many gliomas and glioma cell lines. Transgene replacement causes microtubule disruption and strongly reduces the number of stable clones expressing SIRT2 compared to a control in colony formation assays (Hiratsuka, M et al (2003) Biochem Biophys Res Commun. 309(3) 558-566).

Human SIRT3 is synthesized as an inactive proenzyme and activated by proteolysis on insertion into the mitochondrial matrix. Its function is unknown

5 Human SIRT1 is the closest human homologue to yeast SIR2. It is a nuclear protein found throughout the nucleus. Immunostaining of cells with anti-SIRT1 antibodies shows diffuse nuclear staining.

10 SIRT1 interacts with p53 via the p53 core and carboxy-terminus. It appears to act as a p53 deacetylase, as overexpression of SIRT1 results in reduced acetylation of p53. This in turn leads to reduced expression of endogenous p21, reduced transcription from a p21 reporter construct, and reduced apoptosis in response  
15 to H<sub>2</sub>O<sub>2</sub> and  $\gamma$ -rays.

Overexpression of a catalytically inactive mutant SIRT1 enhances acetylation of p53, and sensitises cells to apoptosis induced by H<sub>2</sub>O<sub>2</sub> and  $\gamma$ -rays.

20 The current consensus is that SIRT1 negatively regulates p53 function via deacetylation of p53, so that inhibition of SIRT1 function sensitises cells to p53-dependent apoptosis in response to cellular stress (Luo et al., 2001; Vaziri et al., 2001;  
25 Langley et al, 2002)

However, modulation of SIRT1 activity has until now been achieved by treatment of cells with the SIRT1 inhibitor nicotinamide, or by overexpression *in trans* of wild type and  
30 catalytically inactive forms of SIRT1. The use of nicotinamide is problematic, partly due to potential inhibition of the SIRT1-related NDACs and SIRT2 and SIRT3, and partly due to potential



pleiotropic effects as nicotinamide is a natural cellular intermediary metabolite. Recent studies have also shown that transgene dosage may be critical in the analysis of p53 function (Blattner *et al.*, 1999; Dumaz *et al.*, 2001), and may underlie the sometimes-conflicting results that have been reported for p53 using this technique.

### Summary of the Invention

In order to elucidate the role of SIRT1 in regulating p53 deacetylation, p53 function and apoptosis, we sought a specific and efficient method of inhibiting the expression or activity of SIRT1.

We addressed the problems of the prior art methods by using a more specific technique, RNA interference, to inhibit SIRT1 expression. RNA interference (RNAi) was used to analyse the function of p53 carboxy-terminal acetylation in p53-dependent apoptosis. The aim was to stabilise acetylation of p53 by silencing of SIRT1. RNAi is a sequence-specific post-translational gene silencing mechanism, which can be initiated in cultured mammalian cells by transfection of a 19-21 nucleotide RNA duplex (short interfering RNA; 'siRNA') homologous in sequence to the target mRNA (Elbashir *et al.*, 2001). Importantly, it has been demonstrated that RNAi does not in itself engage the apoptotic machinery nor alter apoptotic processes, (Jiang and Milner, 2002; Jiang and Milner 2003).

Using RNAi to inhibit SIRT1 expression, we surprisingly found that inhibition of SIRT1 induces massive apoptosis in tumour cells even without additional apoptotic stimuli. This effect is independent of p53 and thus appears to represent a completely

new and unexpected activity of SIRT1. It also appears to be independent of the known pro-apoptotic proteins Bax and PUMA.

Even more surprisingly, the induction of apoptosis seen on inhibition of SIRT1 expression by RNAi appears to be specific to tumour cells, as it does not occur in normal human fibroblasts. This opens the way for new approaches to cancer treatment via the specific induction of apoptosis in tumour cells.

Prior to the invention, inhibition of SIRT1 expression would not have been expected to induce apoptosis in tumour cells in preference to normal cells. In fact, the opposite assumption would have been made, as it was thought that SIRT1 acted only on the p53-dependent pathway of apoptosis, and many tumour cells lack functional p53. Further, it would not have been expected that inhibition of SIRT1 expression alone would induce apoptosis in the absence of other stimuli to trigger p53-dependent apoptotic pathways. The observations of the prior art indicated that inhibition of SIRT1 expression would exacerbate the apoptotic phenotype of cells dying in response to agents that activate p53-dependent apoptotic pathways (Luo et al., 2001; Vaziri et al., 2001), but not that loss of SIRT1 expression itself could induce apoptosis in a p53-independent manner.

The invention accordingly provides for a method of inducing apoptosis in a cell comprising administering a SIRT inhibitor to the cell. Preferably, the cell is a tumour cell. In some embodiments, the cell lacks functional p53, Bax and/or PUMA protein.

The SIRT1 inhibitor may be an agent for inducing RNA interference in a cell, such as a siRNA, a dsRNA, or a nucleic

acid encoding such RNA. In a preferred embodiment, said agent is a siRNA.

5 In another aspect, the invention provides a method of treating a proliferative disease comprising administering to an individual in need thereof an effective amount of a SIRT1 inhibitor. The disease may be cancer, for example a colorectal carcinoma.

10 Thus, the invention also provides a SIRT1 inhibitor for use in a method of medical treatment or therapy. The therapy may be treatment of a proliferative disease, for example cancer. Also provided is the use of a SIRT1 inhibitor in the manufacture of a medicament for the treatment of a proliferative disease.

15 In another aspect, the invention provides a method of identifying a SIRT inhibitor for use in a method of treatment of the invention, the method comprising administering a candidate compound to cultured tumour cells in vitro; determining whether SIRT expression and/or activity is reduced in said cells; and  
20 assaying for apoptosis of said cells. The method may further comprise a step of administering said candidate compound to cultured non-tumour cells in vitro and assaying for apoptosis of said cells.

25 In another aspect, the invention provides an agent for inhibiting the expression of SIRT1 protein in a cell. The agent may be an agent which induces RNA interference to SIRT1 mRNA in a cell, for example siRNA, a dsRNA, or a nucleic acid encoding such RNA. In a preferred embodiment, the agent is a siRNA.

Compositions for pharmaceutical use comprising agents of the invention in combination with a pharmaceutically acceptable excipient are also provided.

- 5 The invention will now be described in more detail with reference to the following figures.

**Brief description of the Figures**

**Figure 1.** siRNAs used in this study.

- 10 SIRT1 siRNA (SEQ ID No. 1)  
Lamin A/C siRNA  
BCR-Abl siRNA

**Figure 2**

- 15 **A.** RT-PCR showing that SIRT siRNA suppresses SIRT-1 mRNA expression.  
**B.** RT-PCR of SIRT-1 in p53<sup>+/+</sup> cells (top) and p53<sup>-/-</sup> cells (bottom) transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated.  
20 **C.** Control: RT-PCR of vimentin in p53<sup>+/+</sup> cells (left) and p53<sup>-/-</sup> cells (right) transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated.  
**D.** Control: RT-PCR of Lamin A/C in p53<sup>+/+</sup> cells (left) and p53<sup>-/-</sup> cells (right) transfected with SIRT1 siRNA, Lamin A/C siRNA  
25 and Bcr-Abl siRNA as indicated.

**Figure 3.** Western blots showing that SIRT1 siRNA suppresses SIRT1 protein expression.

- A.** Western blot of whole cell lysates from cells transfected  
30 with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as

indicated. Blots were probed with an anti-SIRT1 antibody and an anti-Lamin A/C antibody as shown.

**B.** Western blot of the soluble fraction of lysates from p53<sup>+/+</sup> cells (left) and p53<sup>-/-</sup> cells (right) transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. Equal amounts of total protein were loaded in each lane. Blots were probed with an anti-SIRT1 antibody. The bands 2' and 10' indicate 2 minute and 10 minute exposure times respectively

**C.** Western blot of the soluble fraction of lysates from p53<sup>+/+</sup> cells transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. The amount of total protein in each lane was not equalised. Blots were probed with an anti-SIRT1 antibody.

**D.** Western blot of the soluble fraction of lysates from p53<sup>-/-</sup> cells transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. The amount of total protein in each lane was not equalised. Blots were probed with an anti-SIRT1 antibody.

**Figure 4** Western blots showing that SIRT1 siRNA stabilises p53 protein and increases p53 protein expression.

**A.** Western blot of whole cell lysates, soluble and insoluble fractions from cells transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. Blots were probed with an anti-p53 antibody.

**B.** Western blot of whole cell lysates (WCE), soluble (SOL) and insoluble (PEL) fractions of lysates from p53<sup>+/+</sup> cells (top) and p53<sup>-/-</sup> cells (bottom) transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. Blots were probed with an anti-p53 antibody. Recombinant human p53 was run as an antibody control.

C. Western blot of whole cell lysates (WCE), soluble (SOL) and insoluble (PEL) fractions of lysates from p53<sup>+/+</sup> cells transfected with SIRT1 siRNA (top), Lamin A/C siRNA (middle) and Bcr-Abl siRNA (bottom). Blots were probed with an anti-p53 antibody. Recombinant human p53 was run as an antibody control.

**Figure 5** Western blots showing that SIRT1 siRNA induces phosphorylation of p53 protein at serine 15.

A. Western blots of whole cell lysates, soluble and insoluble fractions from cells transfected with SIRT1 siRNA. Blots were probed with the anti-p53 antibody DO-1 (top) and an anti-phosphoSer15 antibody (middle). The bottom blot was probed with antibodies to p21 and HDM2. Cells were harvested at 12, 24, 36 and 48 hours post-transfection, as indicated.

B. Western blot of whole cell lysates (WCE), soluble (SOL) and insoluble (PEL) fractions of lysates from p53<sup>+/+</sup> cells transfected with SIRT1 siRNA (top right), Lamin A/C siRNA (bottom left) and Bcr-Abl siRNA (bottom right). Blots were probed with an anti-phosphoSer15 antibody. Recombinant human p53 was run as an antibody control.

C. Western blot of whole cell lysates (WCE) from cells transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. Blots were probed with an anti-phosphoSer15 antibody.

**Figure 6.** Western blots showing that SIRT1 siRNA induces expression of p21.

A. Western blot of soluble lysates from p53<sup>+/+</sup> cells (left) and p53<sup>-/-</sup> cells (right) transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. Blots were probed with an anti-p21 antibody.

**B.** Western blot of whole cell lysates from p53<sup>+/+</sup> cells incubated at 37°C (bottom) and subjected to cold shock (top) and transfected with SIRT1 siRNA, Lamin A/C siRNA and Bcr-Abl siRNA as indicated. Blots were probed with an anti-p21 antibody.

5 SIRT1 inhibition does not induce p21 expression under conditions of cold shock.

**Figure 7** Controls: Effect of SIRT siRNA on expression of Lamin A/C and Bcr-Abl.

10

**Figure 8** SIRT siRNA induces apoptosis in p53<sup>+/+</sup> tumour cells.

**A** Phase contrast images of HCT116 p53<sup>+/+</sup> cells 24 hours and 48 hours after transfection with SIRT1 siRNA, BCR-Abl siRNA and Lamin A/C siRNA.

15 **B** Phase contrast images of HCT116 p53<sup>+/+</sup> cells at 12 hour timepoints after transfection with SIRT siRNA.

**Figure 9** SIRT siRNA induces apoptosis in p53<sup>+/+</sup> tumour cells.

**A** Phase contrast images of HCT116 p53<sup>-/-</sup> cells 24 hours, 48 hours after transfection with SIRT1 siRNA, BCR-Abl siRNA and Lamin A/C siRNA.

20

**Figure 10** SIRT siRNA induces apoptosis in Bax<sup>-/-</sup> tumour cells.

**A** Phase contrast images of HCT116 p53<sup>+/+</sup> Bax<sup>-/-</sup> cells 24 hours and 48 hours and after transfection with SIRT1 siRNA.

25

**Figure 11** FACS analysis showing apoptosis induced in SIRT1-transfected HCT116 p53<sup>+/+</sup> cells. Cells were analysed at 48 hours post-transfection.

30

**Figure 12** Western blots showing Bax and PUMA expression in cells treated with SIRT1 siRNA. Blots of lysates from cells

transfected with SIRT1 siRNA were probed with an anti-Bax antibody and an anti-PUMA antibody as shown.

**Figure 13** SIRT1 siRNA does not induce apoptosis in primary human fibroblasts.

Phase contrast images of primary human normal diploid fibroblasts (NDFs) transfected with SIRT1 siRNA, taken at 2, 3, 4 and 5 days post-transfection.

**Figure 14** cDNA sequence of human SIRT1 (SEQ ID No. 2).

**Figure 15** Amino acid sequence alignment of human SIRT1 (SEQ ID No. 3), 2 and 3. Asterisks indicate conserved residues. ':' represents conservative amino acid substitution and '.' represents semi-conservative amino acid replacement.

**Figure 16** cDNA sequence alignment of human SIRT1-7 to either side of the region of SIRT 1 covered by the SIRT-1 siRNA (shown in bold) to demonstrate specificity of SIRT1 siRNA sequences..

**Figure 17** Amino acid sequence alignment of human SIRT1-7.

### Detailed description

#### **Proteins**

SIRT1 refers to human sirtuin 1 as described above. SIRT1 is encoded by a nucleic acid with nucleotide sequence shown in SEQ ID No. 2 and Fig. 14. The full-length SIRT1 protein has the sequence shown in SEQ ID No. 3 and Fig. 15. The GenBank accession number for the full-length SIRT1 cDNA and amino acids sequences is NM012238.



Functional p53 protein indicates p53 protein which is capable of inducing apoptosis. A cell lacking functional p53 protein may, for example, carry a deletion or termination mutation of p53 so that it does not express full length p53 at all. Alternatively, the cell may express a mutant p53 which is not capable of triggering apoptosis.

PUMA (p53 upregulated modulator of apoptosis) is a mitochondrial protein that appears to be required for p53-mediated apoptosis (Yu et al., 2003). Expression of PUMA is tightly regulated by p53. PUMA triggers apoptosis via Bax by binding to Bcl-2 and Bcl-X<sub>L</sub>, which otherwise bind and sequester Bax.

Bax is member of the Bcl-2 family of apoptosis regulatory proteins which induces apoptosis, at least in part by triggering release of cytochrome c from mitochondria. Cytochrome c mediates the subsequent activation of the caspases which carry out the apoptotic death programme. Expression of Bax may also be upregulated by p53.

#### ***SIRT1 inhibitor***

The term 'SIRT1 inhibitor' is intended to cover any agent that reduces the expression or activity of SIRT1 in a cell.

Alternatively, the agent may be an agent that inhibits the transcription or translation of SIRT, such as an antisense DNA, RNA or an agent that induces RNA interference.

Inhibition of SIRT expression may be detected by RT-PCR using SIRT1-specific primers, or by Western blotting using an anti-SIRT1 antibody.

### **RNA interference**

RNA interference (RNAi) is a process whereby the introduction of double stranded RNA (dsRNA) into a cell inhibits gene expression post-transcriptionally, in a sequence dependent fashion. This process is also known as post-transcriptional gene silencing. Current models of RNAi indicate that it is mediated by short (typically 20-25 nucleotides) dsRNAs known as 'small interfering RNAs' (siRNA). It appears that dsRNA is cleaved in the cell to create siRNAs. siRNAs are then incorporated into an RNA-induced silencing complex (RISC), guiding the complex to the homologous endogenous mRNA. The activated RISC then cleaves the mRNA transcript, resulting in the destruction of the mRNA in a cell which is homologous to the siRNAs. The siRNAs are re-cycled. In this way, a relatively small number of siRNAs can selectively destroy a large excess of cellular mRNA.

To induce RNA interference in a cell, dsRNA may be introduced into the cell as an isolated nucleic acid fragment or via a transgene, plasmid or virus. Alternatively, siRNA may be synthesised and introduced directly into the cell.

siRNA sequences are selected on the basis of their homology to the gene it is desired to silence. Homology between two nucleotide sequences may be determined using a variety of programs including the BLAST program, of Altschul et al. (1990) *J. Mol. Biol.* 215: 403-10, or BestFit, which is part of the Wisconsin Package, Version 8, September 1994, (Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA, Wisconsin 53711). Sequence comparisons may be made using FASTA and FASTP (see Pearson & Lipman, 1988. *Methods in Enzymology* 183: 63-98). Parameters are preferably set, using the default matrix, as

follows: Gapopen (penalty for the first residue in a gap): -16 for nucleic acid; Gapext (penalty for additional residues in a gap): -4 for nucleic acids; KTUP word length: 6 for nucleic acids.

5

Sequence comparison may be made over the full length of the relevant sequence, or may more preferably be over a contiguous sequence of about or 10, 15, 20, 25 or 30 bases.

- 10 Preferably the degree of homology between the siRNA and the target gene is at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99%.

- 15 The degree of homology between the siRNA or dsRNA and the gene to be silenced will preferably be sufficient that the siRNA or dsRNA will hybridise to the nucleic acid of the gene sequence under stringent hybridisation conditions.

- 20 Typical hybridisation conditions use 4-6 x SSPE; 5-10x Denhardt's solution, 5g polyvinylpyrrolidone and 5g bovine serum albumin; 100µg-1mg/ml sonicated salmon sperm DNA; 0.1-1% sodium dodecyl sulphate; optionally 40-60% deionised formamide. Hybridisation temperature will vary depending on the GC content of the nucleic acid target sequence but will typically be between 42°C-65°C.

- 25 Sambrook et al (2001) Molecular Cloning: A Laboratory Approach (3<sup>rd</sup> Edn, Cold Spring Harbor Laboratory Press). A common formula for calculating the stringency conditions required to achieve hybridisation between nucleic acid molecules of a specified homology is:

- 30  $T_m = 81.5^{\circ}\text{C} + 16.6\text{Log}[\text{Na}^+] + 0.41[\%G+C] - 0.63(\% \text{ formamide}).$

The siRNA may be between 10bp and 30bp in length, preferably

between 20bp and 25bp. Preferably, the siRNA is 19, 20, 21 or 22bp in length.

5 The siRNA sequence may be, for example, any suitable contiguous sequence of 10-30bp from the sequence shown in Fig 14 (SEQ ID No. 2). Alternatively, longer dsRNA fragments comprising contiguous sequences from the sequences of Fig 14 (SEQ ID No. 2) may be used, as they will be cleaved to form siRNAs within the cell. Preferably, the siRNA sequence is that shown in Fig 1a  
10 (SEQ ID No. 1).

15 In a preferred embodiment, the siRNA has the 19bp sequence shown in Fig 1. In some embodiments, the siRNA has an overhang at one or both ends of one or more deoxythymidine bases. The overhang is not to be interpreted as part of the siRNA sequence. Where present, it serves to increase the stability of the siRNA within cells by reducing its susceptibility to degradation by nucleases.

20 siRNA molecules may be synthesized using standard solid or solution phase synthesis techniques which are known in the art. Linkages between nucleotides may be phosphodiester bonds or alternatives, for example, linking groups of the formula  $P(O)S$ , (thioate);  $P(S)S$ , (dithioate);  $P(O)NR'^2$ ;  $P(O)R'$ ;  $P(O)OR_6$ ;  $CO$ ; or  
25  $CONR'^2$  wherein  $R$  is  $H$  (or a salt) or alkyl (1-12C) and  $R_6$  is alkyl (1-9C) is joined to adjacent nucleotides through-O-or-S-.

Alternatively, siRNA molecules or longer dsRNA molecules may be made recombinantly by transcription of a nucleic acid sequence,  
30 preferably contained within a vector as described below.

Modified nucleotide bases can be used in addition to the

naturally occurring bases, and may confer advantageous properties on siRNA molecules containing them.

For example, modified bases may increase the stability of the siRNA molecule, thereby reducing the amount required for silencing. The provision of modified bases may also provide siRNA molecules which are more, or less, stable than unmodified siRNA.

The term 'modified nucleotide base' encompasses nucleotides with a covalently modified base and/or sugar. For example, modified nucleotides include nucleotides having sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group at the 5' position. Thus modified nucleotides may also include 2' substituted sugars such as 2'-O-methyl- ; 2-O-alkyl ; 2-O-allyl ; 2'-S-alkyl; 2'-S-allyl; 2'-fluoro- ; 2'-halo or 2; azido-ribose, carbocyclic sugar analogues  $\alpha$ -anomeric sugars; epimeric sugars such as arabinose, xyloses or lyxoses, pyranose sugars, furanose sugars, and sedoheptulose.

Modified nucleotides are known in the art and include alkylated purines and pyrimidines, acylated purines and pyrimidines, and other heterocycles. These classes of pyrimidines and purines are known in the art and include pseudoisocytosine, N4,N4-ethanocytosine, 8-hydroxy-N6-methyladenine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5 fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyl uracil, dihydrouracil, inosine, N6-isopentyl-adenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-

methyladenine, 7-methylguanine, 5-methylaminomethyl uracil, 5-  
 methoxy amino methyl-2-thiouracil, -D-mannosylqueosine, 5-  
 methoxycarbonylmethyluracil, 5methoxyuracil, 2 methylthio-N6-  
 isopentenyladenine, uracil-5-oxyacetic acid methyl ester,  
 5 psueouracil, 2-thiocytosine, 5-methyl-2 thiouracil, 2-  
 thiouracil, 4-thiouracil, 5methyluracil, N-uracil-5-oxyacetic  
 acid methylester, uracil 5-oxyacetic acid, queosine, 2-  
 thiocytosine, 5-propyluracil, 5-propylcytosine, 5-ethyluracil,  
 5ethylcytosine, 5-butyluracil, 5-pentyluracil, 5-pentylcytosine,  
 10 and 2,6,diaminopurine, methylpsuedouracil, 1-methylguanine, 1-  
 methylcytosine.

### Vectors

The invention also provides vectors comprising a nucleotide  
 15 sequence encoding an siRNA or longer RNA or DNA sequence for  
 production of dsRNA. The vector may be any RNA or DNA vector.  
 The vector is preferably an expression vector, wherein the  
 nucleotide sequence is operably linked to a promoter compatible  
 with the cell. The vector will preferably have at least two  
 20 promoters, one to direct expression of the sense strand and one  
 to direct expression of the antisense strand of the dsRNA.  
 Alternatively, two vectors may be used, one for the sense strand  
 and one for the antisense strand. Alternatively the vector may  
 encode RNAs which form stem-loop structures which are  
 25 subsequently cleaved by the cell to produce dsRNA.

Where the vector is an expression vector, the sequence to be  
 expressed will preferably be operably linked to a promoter  
 functional in the target cells. Promoters suitable for use in  
 30 various vertebrate systems are well known. For example,  
 suitable promoters include viral promoters such as mammalian

retrovirus or DNA virus promoters, e.g. MLV, CMV, RSV, SV40 IEP and adenovirus promoters and metallothionein promoter. The CMV IEP may be more preferable for human use. Strong mammalian promoters may also be suitable as well as RNA polymerase II and  
 5 III promoters. Variants of such promoters retaining substantially similar transcriptional activities may also be used.

Other vehicles suitable for use in delivering nucleic acids such  
 10 as siRNAs include viruses and virus-like particles (VLPs) such as HPV VLPs comprising the L1 and/or L2 HPV viral protein; or hepatitis B viral proteins. Other suitable VLPs may be derived from picornaviruses; togaviruses; rhabdoviruses; orthomyxoviruses; retroviruses; hepadnaviruses; papovaviruses;  
 15 adenoviruses; herpesviruses; and pox viruses.

### ***Delivery***

Various agents may be used to improve the delivery of RNA, DNA or protein into the cell. Viral vectors as described above may  
 20 be used to deliver nucleic acid into a cell. Where other vectors, or no vector, is used, delivery agents such as liposomes may usefully be employed. Delivery peptides such as Antennapedia of the HIV TAT peptide may be used, as may organic polymers such as dendrimers or polylysine-transferrine-  
 25 conjugates.

Liposomes can be prepared from a variety of cationic lipids, including DOTAP, DOTMA, DDAB, L-PE, and the like. Lipid carrier mixtures containing a cationic lipid, such as N-[1-(2,3-  
 30 dioleyloxy) propyl]-N,N,N-triethylammonium chloride (DOTMA) also known as "lipofectin", dimethyl dioctadecyl ammonium bromide

(DDAB), 1,2-dioleoyloxy-3-(trimethylammonio) propane (DOTAP) or L-lysinyphosphatidylethanolamine (L-PE) and a second lipid, such as dioleoylphosphatidylethanolamine (DOPE) or cholesterol (Chol), are particularly useful for use with nucleic acids.

- 5 DOTMA synthesis is described in Felgner, et al., (1987) Proc. Nat. Acad. Sciences, (USA) 84:7413-7417. DOTAP synthesis is described in Stamatatos, et al., Biochemistry, (1988) 27:3917-3925.

- 10 Liposomes are commercially available from many sources. DOTMA:DOPE lipid carriers can be purchased from, for example, BRL. DOTAP:DOPE lipid carriers can be purchased from Boehringer Mannheim. Cholesterol and DDAB are commercially available from Sigma Corporation. DOPE is commercially available from Avanti  
15 Polar Lipids. DDAB:DOPE can be purchased from Promega. Invitrogen make liposomes under the names oligofectamine™ and lipofectamine™.

- To incorporate nucleic acid into liposomes, the liposome-nucleic  
20 acid complex is prepared by mixing with the nucleic acid in an appropriate nucleic acid:lipid ratio (for example 5:3) in a physiologically acceptable diluent (for example Opti-MEM™ at an appropriate dilution) immediately prior to use.

25 ***Apoptosis assays***

The induction of apoptosis in cells may be assayed by many methods.

- Apoptotic cells in culture may be detected and assayed by  
30 photomicrography; apoptotic cells may be detected by their distinctive morphology, with blebbing of the plasma membrane and



chromatin condensation and fragmentation. DNA dyes such as propidium iodide (PI) or Hoechst 33342 may be used to detect chromatin condensation. Alternatively, TUNEL may be used to detect DNA strand breaks.

5

Antibodies such as anti-annexin V can be used to label apoptotic cells and detected by immunofluorescence, and assayed by micrography of FACS analysis.

- 10 FACS analysis may be used in combination with a DNA dye such as PI, TUNEL and/or annexin staining. Apoptotic cells may be detected as a sub-G1 fraction of cells which have lost DNA.

- 15 Release of cytochrome c from mitochondria is another marker for apoptosis. A measure of the early stages of apoptosis may be obtained by fractionating cell lysates into mitochondrial and cytoplasmic fractions and detecting the amount of cytochrome c released from the mitochondria into the cytosol. This is usually done by performing a Western blot and probing with an  
20 anti-cytochrome c antibody.

### ***Proliferative disease***

- A proliferative disease is a pathological condition characterised by unwanted cell growth. In general,  
25 proliferative diseases can be divided into two types: clonal and non-clonal. Clonal proliferative disease usually leads to the formation of tumours, which may be benign or malignant (cancerous). Cancers may be cancer of the skin, connective tissue, adipose, breast, lung, stomach, pancreas, ovary, cervix,  
30 uterus, kidney, bladder, colon, prostate, central nervous system (CNS), retina and circulating tumours (such as leukaemia and

lymphoma). Colorectal cancer includes cancers of the colon, rectum, anus, and appendix.

5 SIRT inhibitors may be effective in providing treatments that discriminate between malignant and normal cells, avoiding many of the deleterious side-effects present with most current chemotherapeutic regimes.

10 Non-clonal proliferative diseases include psoriasis, fibrocystic disease, myelofibrosis, proliferative diabetic retinopathy, atherosclerosis (associated with proliferation of vascular cells) and chronic inflammatory proliferative diseases (CIPD).

15 As used herein, 'tumour cells' shall be taken to refer both to cells derived from tumours, including malignant tumours, and cells immortalised in vitro. 'Normal' cells refers to cells with normal growth characteristics that do not show abnormal proliferation.

20 'Therapy' and 'treatment' of disease includes any therapy or treatment that alleviates in any way the symptoms of a disease. These terms refer to any administration of the compound, salt or N-oxide thereof, intended to alleviate the severity of a disorder of the GI tract in a subject, and includes treatment  
25 intended to cure the disease, provide relief from the symptoms of the disease and to prevent or arrest the development of the disease in an individual at risk from developing the disease or an individual having symptoms indicating the development of the disease in that individual.

### *Compositions*

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may include, in addition to active ingredient, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous or intravenous.

For solid compositions, conventional non-toxic solid carriers include, for example, pharmaceutical grades of mannitol, lactose, cellulose, cellulose derivatives, starch, magnesium stearate, sodium saccharin, talcum, glucose, sucrose, magnesium carbonate, and the like may be used. The active compound as defined above may be formulated as suppositories using, for example, polyalkylene glycols, acetylated triglycerides and the like, as the carrier. Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc, an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline aqueous dextrose, glycerol, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, sorbitan monolaurate, triethanolamine oleate, etc. Actual methods of preparing such

dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, 20th Edition, 2000, pub. Lippincott, Williams & Wilkins. The composition or formulation to be administered will, in any  
5 event, contain a quantity of the active compound(s) in an amount effective to alleviate the symptoms of the subject being treated.

#### **Administration**

10 Where a composition as described herein is to be administered to an individual, administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to  
15 the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. It will also depend upon toxicity of the therapeutic agent, as determined by pre-clinical and clinical trials.

20

Dosing is dependent on severity and responsiveness of the condition to be treated, with course of treatment lasting from several days to several months or until a diminution of disease state is achieved. Optimal dosing schedules are easily  
25 calculated from measurements of drug accumulation in the body. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Therapeutically or prophylactically effective amounts (dosages) may vary depending on the relative potency of individual compositions, and can  
30 generally be routinely calculated based on molecular weight and EC50s in in vitro and/or animal studies. For example, given the molecular weight of an siRNA drug compound (derived from

oligonucleotide sequence and chemical structure) and an experimentally derived effective dose such as an  $IC_{50}$ , for example, a dose in mg/kg is routinely calculated. In general, dosage is from 0.001 $\mu$ g to 100g and may be administered once or several times daily, weekly, monthly or yearly.

Compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated.

Administration may be topical (including ophthalmic, vaginal, rectal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous drip, subcutaneous, intraperitoneal or intramuscular injection, pulmonary administration; e.g., by inhalation or insufflation, or intracranial, e.g., intrathecal or intraventricular, administration.

Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors. A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

The invention is illustrated by the following examples.

#### Experimental procedures

##### **Design of siRNA specific to SIRT1**

A siRNA sequence located adjacent to the conserved sirtuin domain on the carboxy-terminal side was selected on the basis of its selectivity for SIRT1 and its predicted lack of secondary structure.

Bestfit matching to other sirtuin cDNAs was performed and the results were: SIRT2/3/7 - 57.9%

SIRT4/5/6 - 63.2%

5

BLAST searching of GenBank, EMBL, DDBJ and PDB databases was performed and the results were:

Matches 1-8 :Human SIRT1 [some truncated clones].

Match 9: *Streptococcus mutans* sequence [imperfect].

10 Match 10: *Mus musculus* sequence [imperfect].

BLAST searching of the Genbank Human EST database was performed and the results were:

Matches 1-4: Human SIRT1.

15 Matches 5-10: Misc. sequences [imperfect].

These results showed that the selected sequence was likely to be specific for SIRT RNA.

20 The biophysical properties and predicted secondary structure formation of the selected sequence was

T<sub>m</sub>: 46°C

GC: 42.1%

Loop: No

25 Secondary structure: No

Taken together, these results indicated that the selected sequence was likely to be effective in inducing RNA interference specific for the SIRT1 mRNA.

30

Control siRNAs were Lamin A/C siRNA (purchased from Dharmacon) and Bcr-Abl siRNA (siACE RNA, obtained from Dr Ming Jiang). A

further control, indicated in the figures as CTRL or Mock, was the transfection of siRNA buffer alone (no siRNA).

#### RT-PCR

5 RT-PCR for SIRT1, Lamin A/C, BCR-Abl and controls (vimentin and GAPDH) was performed.

A SIRT1 upstream [5'] primer located in conserved core domain, over an exon-exon boundary, and a downstream [3'] primer located in SIRT1-specific sequence were selected.

10 RT-PCR was performed from total RNA using a Reverse-It One-Step kit (ABgene) according to the manufacturer's instructions, and visualised on 2% TAE gels with ethidium bromide.

Primers were as follows:

SIRT1 sense 5-TCAGTGTCATGGTTCCTTTGC-3; SIRT1 antisense 5-

15 AATCTGCTCCTTTG

CCACTCT;

Lamin A sense 5-AAGCAGCGTGAGTTTGAGAGC-3;

Lamin A antisense 5-AGGGTGAACTTTGGTGGGAAC-3;

GAPDH sense 5-CGGAGTCAACGGATTTG

20 GTCGTAT-3;

GAPDH antisense 5-AGCCTTCTCCATGGTGGTGAAGAC-3.

Vimentin sense: 5-gCCAACCTACATCgACAAGgTg-3

Vimentin antisense: 5-gAgCAggTCTTggTATTCACg-3

25

#### Western blotting

Cells for protein lysates were washed in PBS and lysed in buffer IPAX (10 mM TRIS base pH 8.0, 140 mM NaCl, 0.5% NP40, 1 mM PMSF, 1x Complete Protease Inhibitors (Roche), 50 mM NaF, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 30 1 µM trichostatin A, 10 mM nicotinamide). Lysates were spun at 13 000g for 20 minutes at 4°C to separate soluble and insoluble fractions, where indicated. Mitochondrial and cytosolic

fractions were prepared as previously described (Marsden *et al.*, 2002), where indicated. Lysates were run on 10% or 15% SDS-PAGE, electroblotted to Protran membrane (Schleicher & Schuell), and probed with antibodies as described below. Visualisation was with the POD chemiluminescence kit (Roche).

Blots were probed with the following antibodies:

anti-SIRT1 (H-300, Santa Cruz); anti-p53 (DO-1, Oncogene); anti-phosphoserine 15 (Ser15-R, Santa Cruz), anti-p21 (SX118, Pharmingen); anti-HDM2 (monoclonal antibody 4B2, prepared in-house); anti-lamin A/C (636, Santa Cruz); Anti-Abl (8E9, Pharmingen); anti-Bax (N-20, Santa Cruz); anti-PUMA $\alpha$  (AHP727, Serotec) and anti-cytochrome c (7H8.2C12, Pharmingen).

## 15 Photomicrography

Phase contrast images were captured with an Axiovert 200M Cell Observer platform (Zeiss) at various time points post-transfection.

## 20 Cell cycle analysis

Cells were fixed with 70% ethanol at  $-20^{\circ}\text{C}$ , washed with PBS and incubated for 15 minutes at room temperature in PBS containing  $100\text{ }\mu\text{g ml}^{-1}$  RNaseA and  $30\text{ }\mu\text{g ml}^{-1}$  propidium iodide. Samples were analysed on a FACSCalibur flow cytometer (Beckton Dickinson) using CellQuest software.

## Results

### Example 1

#### Inhibition of SIRT expression by RNAi

30 A siRNA specific to human SIRT1 was designed which had no more than ~60% homology to any other human sirtuin cDNA (Figure 1A).



BLAST searching of human genomic and EST databases reported only the SIRT1 gene itself as a significant match. As a positive control we used an siRNA directed to lamin A/C (Dharmacon; Elbashir et al., 2001) (Fig 1B); as a negative control we used an siRNA directed to the BCR-ABL fusion oncogene (MWG-Biotech) which is biologically active only in a background of BCR-ABL (Fig 1C). The siRNAs were introduced to wild-type or p53<sup>-/-</sup> or Bax<sup>-/-</sup> HCT116 colorectal cancer cells by cationic-based lipid transfection (Oligofectamine<sup>TM</sup>, Invitrogen) according to the manufacturer's instructions. Cells were harvested 48 hours post-transfection for the preparation of total cellular RNA and at 12 hour intervals post-transfection for the preparation of protein lysates and protein lysates.

One-step RT-PCR showed that only the SIRT1 siRNA reduced the abundance of SIRT1 mRNA (Fig 2B); the mock transfections and control siRNAs did not alter SIRT1 mRNA. RT-PCR specific for the lamin A mRNA demonstrated that the lamin A/C siRNA was active in HCT116 cells (Fig 2D), and therefore that an active siRNA process does not alter SIRT1 mRNA.

Additionally, silencing of SIRT1 mRNA did not alter lamin A mRNA, and none of the siRNAs altered abundance of the mRNAs chosen as controls, vimentin (Fig 2C) and GAPDH (Fig2A). These data support the specificity of the SIRT1 siRNA for RNAi.

Western blots were performed to establish the efficacy of RNAi in inhibiting protein expression. Figs 3A and 7A/B show that only LaminA/C siRNAs silence Lamin A protein expression in p53<sup>+/+</sup> cells (top) and p53<sup>-/-</sup> cells under optimal growth conditions, but silencing of SIRT1 can downregulate LaminA if cellular stress is applied. (Note that the antibody used does not seem to recognise

Lamin A in whole cell lysates). Fig 7C/D confirms that BCR-Abl siRNA does not induce RNAi in cells lacking BCR-ABL. Fig 3 demonstrates the efficacy of SIRT1 siRNA in reducing SIRT1 protein expression.

5

Significant silencing at the protein level was observed for LaminA/C which had not been indicated by the RT-PCR. This suggests that the relationship between silencing at the mRNA and protein levels is not linear. Likewise, blots probed for SIRT1 showed a profound silencing at the protein level (Figure 3) although RT-PCR had shown only ~50% mRNA degradation. Possibly this reflects protection or sequestration of a pool of the mRNA, such that it is not accessible to components of the RNAi machinery.

15

#### Example 2

##### Effect of SIRT1 siRNA on p53 stability and activity

We went on to investigate the effect of SIRT1 siRNA on p53 stability and activity. In response to cellular stress, it appears that stabilisation of p53 requires only its prior destabilisation by HDM2 (Blattner et al., 1999). The most significant post-translational modification in this regard is phosphorylation of serine 20, now generally accepted as the major mechanism through which p53 is removed from HDM2-mediated negative regulation (Chehab et al., 1999; Dumaz et al., 2001). Phosphorylation of serine 15, previously thought to be involved in stabilisation of p53 (Shieh et al., 1997; Unger et al., 1999), is now believed to be primarily involved in the activation of p53 as a transcription factor (Dumaz and Meek, 1999). The acetylation of the carboxy-terminus of p53 occurs in response to most stimuli that stabilise p53 (Itoh et al., 2001).

Although the function of acetylation of p53 is not fully understood it appears to be stimulated by phosphorylation of serine 15 (Dumaz and Meek, 1999; Lambert *et al.*, 1998), implicated in nucleotide excision repair (Rubbi and Milner, 2003) and in the recruitment of acetylase enzymes to chromatin (Espinosa and Emerson, 2001), and required for efficient p53-dependent apoptosis (Luo *et al.*, 2001; Vaziri *et al.*, 2001). Although it has been demonstrated that, under certain circumstances, neither amino-terminal phosphorylation nor carboxy-terminal acetylation are obligate events for the stabilisation and activation of p53 (Blattner *et al.*, 1999; Ashcroft *et al.*, 2000), the weight of evidence supports a general model whereby stress stimuli signal to p53 via phosphorylation of amino-terminal serine residues. These phosphorylations stabilise p53 and promote interaction with acetylase enzymes; acetylation of the carboxy-terminus fully activates p53 as a transcription factor and may help stabilise p53 (Nakamura *et al.*, 2000).

To determine if silencing of SIRT1 could stabilise p53, we performed Western blots using the pantropic anti-p53 antibody DO-1 in cells treated with SIRT1. This revealed that silencing of SIRT1 led to a significant stabilisation of p53 (Fig 4) compared to the mock transfected control. It is possible that the RNAi process itself constitutes a low-level stress signal, as a small stabilisation of p53 was observed for the lamin A/C siRNA but not the BCR-ABL siRNA. Interestingly, the p53 stabilised by the SIRT1 siRNA was partitioned between the soluble (cytoplasmic and nucleoplasmic) fraction and the insoluble fraction protein, though the p53 stabilised by LaminA/C silencing was found only in the soluble fraction.

We then examined in more detail the stabilisation and activation of p53 in HCT116 cells transfected with SIRT1 siRNA. We chose to examine a four-point time course, with sampling at 12, 24, 36 and 48 hours post-transfection. Protein lysates and fractions  
5 were prepared and blotted with antibodies to p53 and phosphoSer15 (Figs 4 and 5).

Significant stabilisation of p53 was not observed in whole cell lysates until 36 hours post-transfection, although the soluble  
10 fraction showed that there was stabilisation at 24 hours (Fig 4). Intriguingly, stabilised p53 was not detected in the insoluble fraction until 36 hours post-transfection, and was not highly elevated until 48 hours.

15 To analyse the activation of p53 as a transcription factor we examined phosphorylation of serine 15. We determined that stabilised p53 was not subject to serine 15 phosphorylation prior to 36 hours post-transfection. Furthermore, that serine 15  
20 phosphorylation was restricted to p53 in the soluble fraction, as no phosphorylation of this residue was detected at any time point in the insoluble fraction (Fig 5). LaminA/C silencing can induce a very small amount of phosphorylation, and BCR-ABL siRNAs do not stimulate phosphorylation.

25 This data is somewhat counter-intuitive, as one might assume that transcriptionally competent p53 would be associated with chromatin. However, these cells were cultured under normal conditions and in the absence of any exogenous stress stimulus. It was therefore possible that although the p53 was stabilised  
30 and "activated", additional signals or cofactors not present or accessible to the p53 under these conditions were required to activate gene transcription.

Example 3Effect of SIRT1 siRNA on downstream activity of p53

To investigate this, we analysed expression of the two best-characterised p53 responsive genes, p21(CIP1/WAF1) and HDM2. The results for these were essentially identical, as both showed a strong induction of expression at 36 hours that was absent prior to this time point (Figs 5A and 6). There was therefore a good temporal correlation in this data between stabilisation of p53, "activating" post-translational modification, and the induction of transcriptional targets. This effect was seen only in p53<sup>+/+</sup> cells, indicating that the p21 expression was indeed induced by p53 (Fig 6A). Silencing of SIRT1 does not induce expression of p21 if cellular stress is applied (Fig 7B).

Example 4SIRT1 siRNA induces apoptosis in tumour cells independently of p53

As p21 was effectively induced, we wondered if these cells were cell cycle arrested. Phase contrast images (Figs 8 and 9) showed a very clear phenotype that was apparent at 36 hours, and pronounced by 48 hours. Close examination of these cells under higher magnification revealed reduction of the cell bodies, extensive membrane blebbing and a granular appearance to the nucleus (Fig 8C). Surprisingly, this effect was seen in p53<sup>-/-</sup> cells (Fig 9) as well as p53<sup>+/+</sup> cells (Fig 8), indicating that the apoptosis observed was not occurring via a p53-dependent pathway, counter to previous findings.

As these morphological changes are typical of apoptosis, we analysed control and SIRT1-silenced HCT116 cells by flow cytometry (Figure 11). Propidium iodide staining revealed that

there was no significant change in the cell cycle profile of cells transfected with either the lamin A/C or BCR-ABL siRNAs, as compared to the mock-transfected controls. Cells treated with the SIRT1 siRNA showed a marked decrease of the G1 and G2/M populations and a significant increase in sub-G1 population, consistent with apoptotic chromatin fragmentation.

As cells treated with SIRT1 siRNA were clearly undergoing apoptosis, we next considered whether in p53<sup>+/+</sup> cells the stabilised p53 induced the expression of components of the pro-apoptotic machinery. Activation of p53 in HCT116 cells has previously been demonstrated to induce expression of two pro-apoptotic genes, PUMA and Bax (Yu et al., 2003; Zhang et al., 2000). Neither Bax nor PUMA was induced in SIRT1-silenced cells, as compared to the controls (Fig 12A). Indeed, PUMA appeared to be repressed by transfection of the cells with SIRT1 siRNA. Further, SIRT siRNA induced apoptosis even in p53<sup>+/+</sup> cells lacking Bax (Fig 10). These results indicate that SIRT1 siRNA can induce apoptosis via a p53-independent mechanism even in p53<sup>+/+</sup> cells

Two modes of apoptosis are present in human cells: The intrinsic pathway, which functions through mitochondria, and the extrinsic pathway, which functions through the activation of cell death receptors (for review see Green, 1998). PUMA and Bax are both components of the intrinsic pathway, and as neither was induced by the stabilised p53, it was necessary to determine which apoptotic pathway had been activated.

The discovery of cross-talk between the intrinsic and extrinsic pathways (Green, 1998) has been complicated by the discovery that cell type determines the extent of cross-talk. This has led

to the idea that cells fall into one of two types. Type I cells can undergo death receptor induced apoptosis independently of mitochondria; type II cells require mitochondrial involvement for death receptor signalling to induce apoptosis (Scaffidi *et al.*, 1998). Importantly, it has been established that HCT116 cells exhibit the behaviour of type II cells (Deng *et al.*, 2002), which implies that the apoptosis induced by SIRT1 treatment most likely functions through mitochondria.

10 Example 5

SIRT1 siRNA does not induce apoptosis in normal cells

We decided to test the effect of SIRT1 siRNA on a non-tumour cell line. Treatment of primary human normal diploid fibroblasts (NDFs) with SIRT1 siRNA not only failed to provoke the massive apoptosis seen in HCT116 cells, it did not appear to induce apoptosis at all. Cells were still healthy 5 days post-transfection (Fig 13).

We have demonstrated that silencing of SIRT1 in HCT116 colorectal cancer cells provokes massive apoptosis. The tumour suppressor p53 is stabilised in response to SIRT1 silencing, and undergoes at least partial activation for transcriptional function. This effect is independent of p53, and we did not detect induction of either PUMA or Bax, two pro-apoptotic genes that have been shown to be critical for p53-dependent apoptosis in colorectal cancer cells. Moreover, apoptosis was not induced by SIRT1 siRNA in normal cells, indicating this effect may be specific to tumour cells.

### References

- Appella E and Anderson C W. (2001). *Eur. J. Biochem.*, 268, 2764-2772.
- Ashcroft M, Taya Y and Vousden K H. (2000). *Mol. Cell. Biol.*,  
 5 20, 3224-3233.
- Blattner C, Tobiasch E, Liffen M, Rahmsdorf H J and Herrlich P. (1999). *Oncogene*, 18, 1723-1732.
- Chehab N H, Malikzay A, Stavridi E S and Halazonetis T D. (1999). *Proc. Natl. Acad. Sci. USA*, 96, 13777-13782.
- 10 Dumaz N and Meek D W. (1999). *Embo J.*, 18, 7002-7010.
- Dumaz N, Milne D M, Jardine L J and Meek D W. (2001). *Biochem. J.*, 359, 459-464.
- Elbashir S M, Harborth J, Lendeckel W, Yalcin A, Weber K and Tuschl T. (2001). *Nature*, 411, 494-498.
- 15 Espinosa J M and Emerson B M. (2001). *Mol. Cell*, 8, 57-69.
- Frye R. (1999). *Biochem. Biophys. Res. Commun.*, 273, 793-798.
- Green D R. (1998). *Cell*, 94, 695-698.
- Ito A, Lai C-H, Zhao X, Saito S, Hamilton M H, Appella E and Yao, T-P. (2001). *EMBO J.*, 20, 1331-1340.
- 20 Jiang M and Minler J. (2002). *Oncogene*, 21, 6041-6048.
- Jiang M and Milner J. (2003). *Genes Dev.*, 17, 832-837.
- Juan L-J, Shia W J, Chen M H, Yang, W M, Seto E, Lin Y S and Wu C W. (2000). *J. Biol. Chem.*, 275, 20436-20443.
- Lambert P F, Kashanchi F, Radonovich M F, Shiekhattar R and  
 25 Brady J N. (1998). *J. Biol. Chem.*, 273, 33048-33053.
- Langley, E. et al., (2002) *EMBO J* 21/2383-2396.
- Luo J, Nikolaev A Y, Imai S, Chen D, Su F, Shiloh A, Guarente L and Gu W. (2001). *Cell*, 107, 137-148.
- Luo J, Su F, Chen D, Shiloh A and Gu W. (2000). *Nature*, 408,  
 30 377-381.



- Marsden V S, O'Connor L, O'Reilly L A, Silke J, Metcalf D, Ekert P G, Huang D C, Cecconi F, Kuida K, Tomaselli K J, Roy S, Nicholson D W, Vaux D L, Bouillet P, Adams J M and Strasser A. (2002). *Nature*, 419, 634-637.
- 5 Murphy M, Ahn J, Walker K K, Hoffman W H, Evans R M, Levine A J and George D L. (1999). *Genes Dev.*, 13, 2490-2501.
- Nakamura S, Roth J A and Mukhopadhyay T. (2000). *Mol. Cell. Biol.*, 20, 3224-3233.
- Pluquet O and Hainaut P. (2001). *Cancer Lett.*, 174, 1-15.
- 10 Rosenberg M I and Parkhurst S M. (2002). *Cell*, 109, 447-458.
- Rubbi C P and Milner J. (2003). *EMBO J.*, 22, 975-986.
- Scaffidi C, Fulda S, Srinivasan A, Friesen C, Li F, Tomaselli K J, Debatin K-M, Kramer P H and Peter M E. (1998). *EMBO J.*, 17, 1675-1687.
- 15 Shieh S-Y, Ikeda M, Taya C and Prives C. (1997). *Cell*, 91, 325-334.
- Takata T and Ishikawa F. (2002). *Biochem. Biophys. Res. Commun.*, 301, 250-257.
- Unger T, Juven-Gershon T, Moallem E, Berger M, Vogt Sionov R, Lozano G, Oren M and Haupt Y. (1999). *EMBO J.*, 18, 1805-1814.
- 20 Vaziri H, Dessain S K, Ng Eaton E, Imai S, Frye R, Pandita T K, Guarente L and Weinberg R A. (2001). *Cell*, 107, 149-159.
- Yu J, Zhenghe W, Kinzler K W, Vogelstein B and Zhang L. (2003). *Proc. Natl. Acad. Sci. USA*, 100, 1931-1936.
- 25 Zhang L, Yu J, Ho Park B, Kinzler K W and Vogelstein B. (2000). *Science*, 290, 989-992.

CLAIMS

1. A method of treating a proliferative disease comprising administering to an individual in need thereof an effective amount of a SIRT1 inhibitor.
2. A method according to claim 1 wherein the disease is cancer.
3. A method according to claim 2 wherein the cancer is a colorectal carcinoma.
4. An in vitro method of inducing apoptosis in a cell comprising administering a SIRT1 inhibitor to said cell.
5. A method according to claim 4, wherein the cell lacks at least one of functional p53, Bax and PUMA protein.
6. A method according to claim 4 or claim 5 wherein the cell is a tumour cell.
7. A method according to any one of claims 1 to 6, wherein the SIRT1 inhibitor is a siRNA, a dsRNA, a nucleic acid encoding such RNA, or a SIRT1 antisense RNA.
8. A SIRT1 inhibitor for use in a method of medical treatment.
9. A SIRT1 inhibitor for use according to claim 8, wherein said treatment is treatment of a proliferative disease.

10. A SIRT1 inhibitor for use according to claim 8 or claim 9 which is a siRNA, a dsRNA, a nucleic acid encoding such RNA or a SIRT1 antisense RNA.
11. Use of a SIRT1 inhibitor in the manufacture of a medicament for the treatment of a proliferative disease.
12. Use according to claim 11, wherein the proliferative disease is cancer.
13. Use according to claim 12, wherein the cancer is a colorectal carcinoma.
14. Use according to claim 12, wherein the cancer cells lack at least one of functional p53, Bax and protein.
15. Use according to any one of claims 11 to 14, wherein the SIRT1 inhibitor is a siRNA, a dsRNA, a nucleic acid encoding such RNA or a SIRT1 antisense RNA.
16. A siRNA which inhibits expression of SIRT1 in a cell.
17. A siRNA according to claim 16 which comprises a contiguous sequence of 10-30bp from the sequence of SEQ ID No. 2.
18. A siRNA according to claim 17 which is between 19 and 22 bp in length.
19. A siRNA according to claim 18 which is 19bp in length.

20. A siRNA according to claim 19 which has the siRNA sequence of SEQ ID No 1.
21. A composition comprising a siRNA according to any one of claims 16 to 20 and a pharmaceutically acceptable excipient.
22. The method of any one of claims 1 to 6, wherein the SIRT1 inhibitor is a siRNA according to any one of claims 16 to 20.
23. The use of any one of claims 11 to 14, wherein the SIRT1 inhibitor is a siRNA according to any one of claims 16 to 20.
24. A method of identifying a SIRT1 inhibitor for use in a method according to any one of claims 1 to 7, the method comprising
- administering a candidate compound to cultured tumour cells in vitro;
  - determining whether SIRT expression and/or activity is reduced in said cells; and
  - assaying for apoptosis of said cells.
25. A method according to claim 24, wherein the cells lack at least one of functional p53, Bax and protein.
26. A method according to claim 24 or claim 25, further comprising the steps of administering said candidate compound to cultured normal cells in vitro and assaying for apoptosis of said cells.



Figure 1




Figure 2

A








Figure 2

B



C



D










Figure 3


A



B



C



D







Figure 4

A



B







Figure 4

C






Figure 5









Figure 5

B



C







Figure 6

A



B







Figure 7

A



B











Figure 7

C



D








24

CTRL



48






Figure 8A

Bcr-Abl





LaminA/C





Figure 8A

SIRT1








Figure 8B









Bcr-Abl

Figure 9



CTRL



24



48



72









Figure 9

SIRT1



LaminA/C



24



48




72



Figure 10

CTRL

24



48



SIRT1







Figure 11



- 1 - Mock
- 2 - SIRT1 siRNA
- 3 - Mock
- 4 - LaminA/C siRNA
- 5 - Mock
- 6 - BCR-ABL siRNA



Figure 12










Figure 13

NDF: Control [Mock] Transfection


72 hours



120 hours



48 hours



96 hours






Figure 13

NDF: SIRT1 siRNA Transfection






Figure 14 (1)

1 ATGGCGGACG AGGCGGCCCT CGCCCTTCAG CCCGGCGGCT CCCCTCGGC  
 TACCGCCTGC TCCGCCGGGA GCGGGAAGTC GGGCCGCCGA GGGGGAGCCG  
 51 GGCGGGGGCC GACAGGGAGG CCGCGTCGTC CCCCGCCGGG GAGCCGCTCC  
 CCGCCCCCGG CTGTCCCTCC GCGCGAGCAG GGGGCGGCCC CTCGGCGAGG  
 101 GCAAGAGGCC GCGGAGAGAT GGTCCCGGCC TCGAGCGGAG CCCGGGCGAG  
 CGTTCTCCGG CGCCTCTCTA CCAGGGGCGG AGCTCGCCTC GGGCCCGCTC  
 151 CCCGGTGGGG CGGCCCCAGA GCGTGAGGTG CCGGCGGGGG CCAGGGGCTG  
 GGGCCACCCC GCCGGGGTCT CGCACTCCAC GGCCGCCGCC GGTCCCCGAC  
 201 CCCGGGTGCG GCGGCGGCGG CGCTGTGGCG GGAGGCGGAG GCAGAGGCGG  
 GGGCCCACGC CGCCGCCGCC GCGACACCGC CCTCCGCCTC CGTCTCCGCC  
 251 CGGCGGCAGG CGGGGAGCAA GAGGCCCAGG CGACTGCGGC GGCTGGGGAA  
 GCCGCCGTCC GCCCTCGTT CTCCGGGTCC GCTGACGCCG CCGACCCCTT  
 301 GGAGACAATG GGCCGGGCCT GCAGGGCCCA TCTCGGGAGC CACCGCTGGC  
 CCTCTGTTAC CCGGCCCGGA CGTCCCGGGT AGAGCCCTCG GTGGCGACCG  
 351 CGACAACTTG TACGACGAAG ACGACGACGA CGAGGGCGAG GAGGAGGAAG  
 GCTGTTGAAC ATGCTGCTTC TGCTGCTGCT GCTCCCGCTC CTCCTCCTTC  
 401 AGGCGGCGGC GGCGGCGATT GGGTACCGAG ATAACCTTCT GTTCGGTGAT  
 TCCGCCGCCG CCGCCGCTAA CCCATGGCTC TATTGGAAGA CAAGCCACTA  
 451 GAAATTATCA CTAATGGTTT TCATTCTGT GAAAGTGATG AGGAGGATAG  
 CTTTAATAGT GATTACCAAA AGTAAGGACA CTTTCACTAC TCCTCCTATC  
 501 AGCCTCACAT GCAAGCTCTA GTGACTGGAC TCCAAGGCCA CGGATAGGTC  
 TCGGAGTGTA CGTTCGAGAT CACTGACCTG AGGTTCCGGT GCCTATCCAG  
 551 CATATACTTT TGTTCAACAA CATCTTATGA TTGGCACAGA TCCTCGAACA  
 GTATATGAAA ACAAGTCGTT GTAGAATACT AACCCTGTCT AGGAGCTTGT  
 601 ATTCTTAAAG ATTTATTGCC GGAAACAATA CCTCCACCTG AGTTGGATGA  
 TAAGAATTC TAAATAACGG CTTTGTATAT GGAGGTGGAC TCAACCTACT  
 651 TATGACACTG TGGCAGATTG TTATTAATAT CCTTTCAGAA CCACCAAAAA  
 AACTGTGAC ACCGTCTAAC AATAATTATA GGAAAGTCTT GGTGGTTTTT  
 701 GGAAAAAAG AAAAGATATT AATACAATTG AAGATGCTGT GAAATTACTG  
 CCTTTTTTTC TTTTCTATAA TTATGTTAAC TTCTACGACA CTTTAATGAC  
 751 CAAGAGTGCA AAAAAATTAT AGTTCTAACT GGAGCTGGGG TGTCTGTTTC  
 GTTCTCACGT TTTTTTAATA TCAAGATTGA CCTCGACCCC ACAGACAAAG  
 801 ATGTGGAATA CCTGACTTCA GGTCAAGGGA TGGTATTTAT GCTCGCCTTG  
 TACACCTTAT GGAAGTGAAGT CCAGTTCCTT ACCATAAATA CGAGCGGAAC



Figure 14 (2)

851 CTGTAGACTT CCCAGATCTT CCAGATCCTC AAGCGATGTT TGATATTGAA  
GACATCTGAA GGGTCTAGAA GGTCTAGGAG TTCGCTACAA ACTATAACTT

901 TATTTTCAGAA AAGATCCAAG ACCATTCTTC AAGTTTGCAA AGGAAATATA  
ATAAAGTCTT TTCTAGGTTC TGGTAAGAAG TTCAAACGTT TCCTTTTATAT

951 TCCTGGACAA TTCCAGCCAT CTCTCTGTCA CAAATTCATA GCCTTGTCAG  
AGGACCTGTT AAGGTCGGTA GAGAGACAGT GTTTAAGTAT CGGAACAGTC

1001 ATAAGGAAGG AAAACTACTT CGCAACTATA CCCAGAACAT AGACACGCTG  
TATTCCTTCC TTTTGATGAA GCGTTGATAT GGGTCTTGTA TCTGTGCGAC

2T1PHO

1051 GAACAGGTTG CGGGAATCCA AAGGATAATT CAGTGTCATG GTTCCTTTGC  
CTTGTCACAC GCCCTTAGGT TTCCTATTAA GTCACAGTAC\*CAAGGAAACG

1101 AACAGCATCT TGCCTGATTT GTAAATACAA AGTTGACTGT GAAGCTGTAC  
TTGTCGTAGA ACGGACTAAA CATTTATGTT TCAACTGACA CTTTCGACATG

1151 GAGGAGATAT TTTTAATCAG GTAGTTCCTC GATGTCCTAG GTGCCCAGCT  
CTCCTCTATA AAAATTAGTC CATCAAGGAG CTACAGGATC CACGGGTCTGA

1201 GATGAACCGC TTGCTATCAT GAAACCAGAG ATTGTGTTTT TTGGTGAAAA  
CTACTTGGCG AACGATAGTA CTTTGGTCTC TAACACAAAA AACCACTTTT

1251 TTTACCAGAA CAGTTTCATA GAGCCATGAA GATGACAAA GATGAAGTTG  
AAATGGTCTT GTCAAAGTAT CTCGGTACTT CATACTGTTT CTACTTCAAC

1301 ACCTCCTCAT TGTTATTGGG TCTTCCCTCA AAGTAAGACC AGTAGCACTA  
TGGAGGAGTA ACAATAACCC AGAAGGGAGT TTCATTCTGG TCATCGTGAT

1351 ATTCCAAGTT CCATACCCCA TGAAGTGCCT CAGATATTAA TTAATAGAGA  
TAAGGTTCAA GGTATGGGGT ACTTCACGGA GTCTATAATT AATTATCTCT

1401 ACCTTTGCCT CATCTGCATT TTGATGTAGA GCTTCTTGGA GACTGTGATG  
TGGAAACGGA GTAGACGTAA AACTACATCT CGAAGAACCT CTGACACTAC

~~~~~  
ACUU

dTdTUGAA

rJF3

1501 UGCUGUAACC CUGUAdTdT  
TGCTGTAACC CTGTAAAGCT TTCAGAAATT ACTGAAAAAC CTCCACGAAC  
ACGACATTGG GACATTTCTGA AAGTCTTTAA TGACTTTTTG GAGGTGCTTG  
ACGACAUUGG GACAU

rJF4





Figure 14 (3)

1551 ACAAAAAGAA TTGGCTTATT TGTCAGAGTT GCCACCCACA CCTCTTCATG  
 TGTTTTTCTT AACCGAATAA ACAGTCTCAA CGGTGGGTGT GGAGAAGTAC  
 1601 TTTCAGAAGA CTCAAGTTCA CCAGAAAGAA CTTCAACCACC AGATTCTTCA  
 AAAGTCTTCT GAGTTCAAGT GGTCTTTCTT GAAGTGGTGG TCTAAGAAGT  
 1651 GTGATTGTCA CACTTTTAGA CCAAGCAGCT AAGAGTAATG ATGATTTAGA  
 CACTAACAGT GTGAAAATCT GGTTTCGTCGA TTCTCATTAC TACTAAATCT  
 1701 TGTGTCTGAA TCAAAAGGTT GTATGGAAGA AAAACCACAG GAAGTACAAA  
 ACACAGACTT AGTTTTCCAA CATACCTTCT TTTTGGTGTG CTCATGTTT  
 1751 CTTCTAGGAA TGTTGAAAGT ATTGCTGAAC AGATGGAAAA TCCGGATTTG  
 GAAGATCCTT ACAACTTTCA TAACGACTTG TCTACCTTTT AGGCCTAAAC  
 1801 AAGAATGTTG GTTCTAGTAC TGGGGAGAAA AATGAAAGAA CTTCAGTGGC  
 TTCTTACAAC CAAGATCATG ACCCCTCTTT TTACTTTCTT GAAGTCACCG  
 1851 TGGAACAGTG AGAAAATGCT GGCCTAATAG AGTGGCAAAG GAGCAGATTA  
 ACCTTGTCAC TCTTTTACGA CCGGATTATC TCACCGTTTC CTCGTCTAAT  
 TC TCACCGTTTC CTCGTCTAA  
 2T1REV  
 1901 GTAGGCGGCT TGATGGTAAT CAGTATCTGT TTTTGCCACC AAATCGTTAC  
 CATCCGCCGA ACTACCATTA GTCATAGACA AAAACGGTGG TTTAGCAATG  
 1951 ATTTTCCATG GCGCTGAGGT ATATTCAGAC TCTGAAGATG ACGTCTTATC  
 TAAAAGGTAC CGCGACTCCA TATAAGTCTG AGACTTCTAC TGCAGAATAG  
 2001 CTCTAGTTCT TGTGGCAGTA ACAGTGATAG TGGGACATGC CAGAGTCCAA  
 GAGATCAAGA ACACCGTCAT TGTCACATC ACCCTGTACG GTCTCAGGTT  
 2051 GTTTAGAAGA ACCCATGGAG GATGAAAGTG AAATTGAAGA ATTCTACAAT  
 CAAATCTTCT TGGGTACCTC CTACTTTCAC TTAACTTCT TAAGATGTTA  
 2101 GGCTTAGAAG ATGAGCCTGA TGTTCCAGAG AGAGCTGGAG GAGCTGGATT  
 CCGAATCTTC TACTCGGACT ACAAGGTCTC TCTCGACCTC CTCGACCTAA  
 2151 TGGGACTGAT GGAGATGATC AAGAGGCAAT TAATGAAGCT ATATCTGTGA  
 ACCCTGACTA CCTCTACTAG TTCTCCGTTA ATTACTTCGA TATAGACACT  
 2201 AACAGGAAGT AACAGACATG AACTATCCAT CAAACAAATC A  
 TTGTCCTTCA TTGTCTGTAC TTGATAGGTA GTTTGTTTGA T



Figure 15

```

hsIRT2 -----
hsIRT3 -----
hsIRT1 MADEAALALQPGGSPSAAGADREAASSPAGEPLAKRPRRDGPGLERSPGEPGGAAPEREV 60

hsIRT2 -----
hsIRT3 -MAFWGWRAAAALRLWGRVVERVEAGGGVGFQACGCR----- 37
hsIRT1 PAAARGCPGAAAAALWREAEEAAAAGGEQEAQATAAAGEGONGPGLQGPSREPLADNL 120

hsIRT2 -----
hsIRT3 -----MAEP-DPSHPLETQAG- 15
hsIRT1 -----LVLGGRDDVSAGLRGSHGARGEPLDAPRPLQRPPRP 73
YDEDDDDGEEEEEEAAAAAIGYRDNLLFGDEIITNGFHSCESEEDRASHASSSDWTPRP 180
: . . . .

hsIRT2 -----KVQEAQSDSDSEG-----GAAGGEADMDFLRNLFSTLSLGSQKERLLDE 61
hsIRT3 -----EVPRAFRRQPRAAAPSEFFSSIKGRRSISFSVGASSVVGSGGSSDK--GK 122
hsIRT1 RIGPYTFVQQHLMIGTDPRTILKDLLPETIPPELDDMTLWQIVINILSEPPKRKRKDI 240
: . . . . * . . .

hsIRT2 LTLEGVARYMQSERCRRVICLVGAGISTSAGIPDFRSPSTGLYD--NLEKYHLPYPEAIF 119
hsIRT3 LSLQDVAELIRARACQRVVMVGAGISTPSGIPDFRSPGSGLYS--NLQQYDLPYPEAIF 180
hsIRT1 NTIEDAVKLLQ--ECKKIIVLTGAGVSVSCGIPDFRS-RDGIYARLAVDFPDLPDPQAMF 297
: . . . . : * : : : . . . . . * : : : . . . . .

hsIRT2 EISYFKKHPEPFFALAKELYPGQFKPTICHYFMRLKDKGLLLRCTQNIOTLERIAGLE 179
hsIRT3 ELPFFFHNPKEPFTLAKELYPGNYKPNVTHYFLRLLHDKGLLLRCTQNIOTLERVSGIP 240
hsIRT1 DIEYFRKDPRPFFKFAKEIYPGQFQPSLCHKFIALSDKEGKLLRNYTQNIOTLEQVAGIQ 357
: : * : . . . . . * : : : . . . . . * : : : . . . . .

hsIRT2 QEDLVEAHGTFTYTSHCVSASCRHEYPLSWMKEKIFSEVTPKCEDCQS-----LVKPDIVF 234
hsIRT3 ASKLVEAHGTFFASATCT--VCQRPFPGEDIRADVMADRVPRCPVCTG-----VVKPDIVF 293
hsIRT1 R--IIQCHGSFATASCL--ICKYKVDCEAVRGDIFNQVPRCPRCPADEPLAIMKPEIVF 413
: : . . . . : * : : : . . . . . * : : : . . . . .

hsIRT2 FGESLPARFFSCMQSDFLKVDLLLVMGTSLQVQPFASLISKAPLSTPRLLINKE----- 288
hsIRT3 FGEPLPQRFLLHV--DFPMADLLLILGTSLEVEPFFASLTEAVRSSVPRLLINRD----- 346
hsIRT1 FGENLPEQFHRAMKYDKDEVDLLVIGSSSLKVRPVALIPSSIPHEVPQILINREPLPHLH 473
*** ** : * : * . . . . . * : : : . . . . .

hsIRT2 -----KAGQSDPFLGMIMGLGGGMDFDSKKAYRDVAWLGECDQG--- 327
hsIRT3 -----LVGP-----LAWHPR--SRDVAQLGDVVHG--- 369
hsIRT1 FDVELLGDGCDVIINELCHRLGGEYAKLCCNPVKLSEITEKPPRTQKELAYLSELPPTPLH 533
* : . . . . : : * : :

hsIRT2 -----CLALAEELLGWKKELEDLVRREHASIDAQSGAGVNPSTSPASP 369
hsIRT3 -----VESLVELLGWTEEMRDVQRETGKLDG-----P 397
hsIRT1 VSEDSSSPERTSPPDSSVIVTLLDQAAKSNDLDVSESKGCMEEKPQEVQTSRNVESIAE 593
: * : . . . . : * : : :

hsIRT2 KKSPPPAKDEARTTEREKPO----- 389
hsIRT3 DK----- 399
hsIRT1 QMENPDLKNVGSSTGEKNERTSVAGTVRKCPWPNRVAKEQISRRLDGNQYLFPPNRYIFH 653

hsIRT1 GAEVYSDSEDDVLSSSSCGSNSDSGTCQSPSLEEPMEDESEIEEFYNGLEDEPDVPERAG 713

hsIRT1 GAGFGTDGDDQEAINEAISVKQEVTDNMNPSNKS 747

```



Figure 16

|                            |      |             |            |            |             |            |
|----------------------------|------|-------------|------------|------------|-------------|------------|
| SIRT1                      | 1426 | -----       | ---GTA---- | -----      | -----       | -----      |
| SIRT2                      | 981  | ctgcct----  | -----      | -----      | -----       | -----      |
| SIRT3                      | 1107 | cgtgggaaagc | ctaGTG---- | -----      | -----       | -----      |
| SIRT4                      | 817  | -----       | -----      | -----      | -----       | -----      |
| SIRT5                      | 821  | --TTAACACG  | GAGACCACCC | CAGctacgaa | cagattcagg  | tttcatttcc |
| SIRT6                      | 934  | -----       | -----      | -----      | -----       | -----      |
| SIRT7                      | 1125 | gcTTAGCTCG  | GCCCCCATCC | TAGggg---- | -----       | -----      |
| <p>***** ***** ***</p>     |      |             |            |            |             |            |
|                            |      |             |            |            |             |            |
| SIRT1                      | 1429 | -----       | -----      | -----GAGC  | TTCTTGGAgA  | ctgtgatgtc |
| SIRT2                      | 987  | -----       | -----GGCC  | CTTGCTGAGC | TCCTTGGATG  | GA-----    |
| SIRT3                      | 1123 | -----       | -----      | -----GAGC  | TTCTGGGCTG  | GA-----    |
| SIRT4                      | 817  | -----       | -----      | -----C     | TCACTGCCTG  | GG-----    |
| SIRT5                      | 869  | agggaccctg  | tggaacGACT | CTTCCTGAAG | CCCTTGCCCTG | tc-----    |
| SIRT6                      | 934  | -----       | -----      | -----C     | CCGCCGGCCC  | CA-----    |
| SIRT7                      | 1151 | -----       | -----      | -----      | -----       | -----      |
| <p>**** ***** ***** **</p> |      |             |            |            |             |            |
|                            |      |             |            |            |             |            |
| SIRT1                      | 1453 | ataattaatg  | aattgtgtca | taggttaggt | ggtgaatatg  | ccaaactttg |
| SIRT2                      | 1013 | -----       | -----      | -----      | -----       | -----      |
| SIRT3                      | 1139 | -----       | -----      | -----      | -----       | -----      |
| SIRT4                      | 830  | -----       | -----      | -----      | -----       | -----      |
| SIRT5                      | 911  | -----       | -----      | -----      | -----       | -----      |
| SIRT6                      | 947  | -----       | -----      | -----      | -----       | -----      |
| SIRT7                      | 1151 | -----       | -----      | -----      | -----       | -----      |
|                            |      |             |            |            |             |            |
| SIRT1                      | 1503 | ctgtaaccct  | gtaaagcttt | cagaaattac | tgaaaaacct  | ccacgaacac |
| SIRT2                      | 1013 | -----       | -----      | -----      | -----       | -----      |
| SIRT3                      | 1139 | -----       | -----      | -----      | -----       | -----      |
| SIRT4                      | 830  | -----       | -----      | -----      | -----       | -----      |
| SIRT5                      | 911  | -----       | -----      | -----      | -----       | -----      |
| SIRT6                      | 947  | -----       | -----      | -----      | -----       | -----      |
| SIRT7                      | 1151 | -----       | -----      | -----      | -----       | -----      |
|                            |      |             |            |            |             |            |
| SIRT1                      | 1553 | aaaaagaatt  | ggcttatttg | tcagagttgc | caccacacac  | tcttcatggt |
| SIRT2                      | 1013 | -----       | -----      | -----      | -----       | -----      |
| SIRT3                      | 1139 | -----       | -----      | -----      | -----       | -----      |
| SIRT4                      | 830  | -----       | -----      | -----      | -----       | -----      |
| SIRT5                      | 911  | -----       | -----      | -----      | -----       | -----      |
| SIRT6                      | 947  | -----       | -----      | -----      | -----       | -----      |
| SIRT7                      | 1151 | -----       | -----      | -----      | -----       | -----      |



Figure 17 (1)

|        |                                                                 |     |
|--------|-----------------------------------------------------------------|-----|
| hSIRT2 | -----MAFWGWRAAAALRLWGRVVERVEAGGGVGPFFQACGCRLVLGGRDDV            | 46  |
| hSIRT3 | MADEAALALQPGGSPSAAGADREAASSPAGEPLRKRPRRDGPGLERSPGEPGGAAPEREV    | 60  |
| hSIRT1 | -----MAAGGLSRSERKAAERVRLREEQQ                                   | 25  |
| hSIRT4 | -----                                                           |     |
| hSIRT7 | -----                                                           |     |
| hSIRT6 | -----                                                           |     |
| hSIRT5 | -----                                                           |     |
| hSIRT2 | -----MAEP-DPSHPLETQAG-KVQEAQDSDSSEGE-----GAAGGEADMDFLR          | 42  |
| hSIRT3 | SAGLRGSHGARGEPLDPAARPLQRPPEVPRAFRRQPRAAAPSFFFSSIKGGRRSISFSV     | 106 |
| hSIRT1 | PAAARGCPGAAAAALWREAEAEAAAAGGEQEAQATAAAGEGDN--GPGLOGPSREPPLAD    | 118 |
| hSIRT4 | -----MKMSFALTFRSAKGRWIANPSQP-----CSKASIGLFV                     | 33  |
| hSIRT7 | RERLR-----QVSRILRKAASAEGRLLAESADLVTELQG-----RSRRREGLKR          | 73  |
| hSIRT6 | -----MSVNYAAGLSPYADK-----GKCGLPPEIFD                            | 25  |
| hSIRT5 | -----MRPLQIVPSRLISQLYCGLKPP-----ASTRNQICLK                      | 32  |
| hSIRT2 | NLFSQTLSLGSQKER-----                                            | 57  |
| hSIRT3 | GASSVVGSGGSSDK-----                                             | 120 |
| hSIRT1 | NLYDEDDDDDEGESEEEEEAAAAAIGYRDNLLFGDEIITNGFHSCESDEEDRASHASSSDWTP | 178 |
| hSIRT4 | PASPP-----                                                      | 38  |
| hSIRT7 | RQEEVCD-----                                                    | 80  |
| hSIRT6 | PPEE-----                                                       | 29  |
| hSIRT5 | MARP-----                                                       | 36  |
| hSIRT2 | -----LLDELTLEGVARYMQSE-----                                     | 74  |
| hSIRT3 | -----GKLSLQDVAELIRAR-----                                       | 135 |
| hSIRT1 | RPRIGPYTFVQQHLMIGTDPRTILKDLLPETIPPELDDMTLWQIVINILSEPPKRKKRK     | 238 |
| hSIRT4 | -----LDPEKVKELQRFITLS-----                                      | 54  |
| hSIRT7 | -----DPEELRGKVRELASAVR-----                                     | 97  |
| hSIRT6 | -----LERKVVWELARLVWQS-----                                      | 44  |
| hSIRT5 | -----SSSMADFRKFFAKA-----                                        | 50  |
| hSIRT2 | -----RCRRVICLVGAGISTSAGIPDFRSPSTGLYD--NLEKYHLPYPEAIF            | 119 |
| hSIRT3 | -----ACQVVVMVGAGISTPSGIPDFRSPGSGLYS--NLQYDLPYPEAIF              | 180 |
| hSIRT1 | DINTIEDAVKLLQECKKIIIVLTGAGVSVSCGIPDFRS--RDGIYARLAVDFPDLPDQAMF   | 297 |
| hSIRT4 | -----KRLLVMTGAGISTESGIPDYRSEKVGLYAR--TDRRPIQHGDVFR              | 97  |
| hSIRT7 | -----NAKYLVVYTGAGISTAASIPDYRG--PNGVWT-----LLQKGRSVS             | 136 |
| hSIRT6 | -----SSVVFHTGAGISTASGIPDFRG--PHGVWT-----MEERGLAP                | 80  |
| hSIRT5 | -----KHIVIISGAGVSAESGVPTFRG--AGGYWR--KWQAQDLATPLAFA             | 92  |
|        | :: **:* . . . : * . * :                                         |     |
| hSIRT2 | EISYFKKHPEPFFALAKELYPGQFKPTICHYFMRLKDKGLLLRCYTQNIIDTLERAGLE     | 179 |
| hSIRT3 | ELPFFFFHNPFPFFTLAKELYPGNYKPNVTHYFLRLLHDKGLLLRLYTQNIIDGLERVSGIP  | 240 |
| hSIRT1 | DIEYFRKDPFPFFKFAKEIYPGQFQPSLCHKFIALSDKEGKLLRNYTQNIIDTLEQVAGIQ   | 357 |
| hSIRT4 | SAPIRQRYWARNEFVGWPQFSS--HQPNPAHWALSTWEKLGKLYWLVTONVDALHTKAGSR   | 155 |
| hSIRT7 | AADLS-----EAEPTLTHMSITRLHEQKLVQHVVSQNCGLHLRSLGLP                | 179 |
| hSIRT6 | KFDTTFES-----ARPTQTHMALVQLERVGLLRFLVSVQNVQDGLHVRSGFP            | 125 |
| hSIRT5 | HNPSRVWEFYHYRREVMSKEPNAGHRAIAECETRLGKQGRVVVITQNIIDELHRKAGTK     | 152 |
|        | ::** * . . . *                                                  |     |





Figure 17 (2)

|        |                                                                 |     |
|--------|-----------------------------------------------------------------|-----|
| hsIRT2 | QEDLVEAHGTFYTSHCVSASCRHEYPLSW-MKEKIFSEVTPKCEDCQS-----LVKP       | 230 |
| hsIRT3 | ASKLVEAHGTFASATCT--VCQRPFPGED-IRADVMADRVPRCPVCTG-----VVKP       | 289 |
| hsIRT1 | R--IIQCHGSFATASCL--ICKYKVDCEA-VRGDI FNQVVPRCPRCPADEP---LAIMKP   | 409 |
| hsIRT4 | R--LTELHGCM DRVLC LD--CGEQTPRGV-LQERFQVLNPTWSAEAHG-----LAPD     | 202 |
| hsIRT7 | RTAISELHGNMYIEVCTSCVPNREYVRVFDV TERTALHRHQTGR TCHKCG-----TQLRD  | 234 |
| hsIRT6 | RDKLAE LHGNMFVEECAK--CKTQYVRD TVVGT MGLKATGRLCTVAKARGLRACRGELRD | 183 |
| hsIRT5 | N--LLEIHGSLFKTRCTS-----CGVVAENYKSPICPALSG-----KGAP              | 190 |
|        | : : * * : *                                                     |     |
| hsIRT2 | DIVFFGES-LPARFFS--CMQSDFLKVDLLVMGTSLQVQ---PFASLISKAPLSTPRLL     | 284 |
| hsIRT3 | DIVFFGEP-LPQRFLL--HVV-DFPMADLLLI LGTSLEVE---PFASLTEAVRSSVPRLL   | 342 |
| hsIRT1 | EIVFFGEN-LPEQFHR--AMKYDKDEV DLLIVIGSSLKVR---PVALIPSSIPHEVPQIL   | 463 |
| hsIRT4 | GDVFLSEE-QVRSFQVPTCVQC GGHLKPDVFFGDTVNP D---KVD FVHKRVKEADSLLV  | 258 |
| hsIRT7 | TIVHFGERTLGQPLNWEAATEAASRADTILCLGSSLKVLKKYPRLWCMTKPPSRPKLY      | 294 |
| hsIRT6 | TILDWEDS-LPDRDLA--LADEASRNADLSITLGTSLQIR---PSGNLPLATKRRGGRLV    | 237 |
| hsIRT5 | EPGTQDASIPVEKLPRCEEAGCGLLRPHVWVWFGENLOPA---ILEEVORELAHCOLCLV    | 247 |
|        | : : * : :                                                       |     |
| hsIRT2 | INKE-----KAGQSDPFLGMIMGLGGGMDFD SKKAYRDVAW                      | 320 |
| hsIRT3 | INRD-----LVGP-----LAWHPR--SRDVAQ                                | 362 |
| hsIRT1 | INREPLPHLHFDVELLGDCDVI INELCHRLGGEYAKLCCNPVKLSEITEKPPRTQKELAY   | 523 |
| hsIRT4 | VGSS-----LQVYSGYR FILTAW EKKLPIAILN                             | 286 |
| hsIRT7 | IVNLQWTP-----KDDWAALKLHGKCD DVMRLMAELGLEIPA                     | 332 |
| hsIRT6 | IVNLQP-----TKHDRHADLRIHG YVDEVMTRLMEHLGLEIPA                    | 275 |
| hsIRT5 | VGTS-----SVVYPAAMFAPQVAARGVPVAE                                 | 273 |
|        | : : * : :                                                       |     |
| hsIRT2 | LGECDOG-----CLALAE LLGWKKELEDLVRREHASIDAQSGAG                   | 359 |
| hsIRT3 | LGDVVHG-----VESLVELLGWTEEMRDLVQRETGKLDG-----                    | 396 |
| hsIRT1 | LSELPTPLHVSEDSSSPERTSPPDSSVIVTLLDQA AKSNDDLDVSESKGCME EK PQEVQ  | 583 |
| hsIRT4 | IGPTRSD-----DLACLKLN SRCGELLPLIDPC-----                         | 314 |
| hsIRT7 | YSRWQDP-----IFSLATPLRAGEEGSHSRKSLCRSREEAPPG                     | 370 |
| hsIRT6 | WDGPRVLER-----ALPPLPRPPTPKLEPKESPTRINGSIPAGPKQEP CA             | 321 |
| hsIRT5 | ENTETTP-----ATNRFRFH FQGPCGTTLPEALACHENETVS-                    | 310 |
|        | : : * : :                                                       |     |
| hsIRT2 | VPNPSTSASPKKSPPPAKDEARTTEREKPO-----                             | 389 |
| hsIRT3 | -----PDK-----                                                   | 399 |
| hsIRT1 | TSRNVESIAEQMENPDLKNVGSSTGEKNERTSVAGTVRK CWPNRVAKEQISRRLDGNQYL   | 643 |
| hsIRT4 | -----                                                           |     |
| hsIRT7 | DRGAPLSSAPILGGWFGRGCTKRTRKKVT-----                              | 400 |
| hsIRT6 | QHNGSEPASPKRERPTSPAPHRPPKRVKAKAVPS-----                         | 355 |
| hsIRT5 | -----                                                           |     |
| hsIRT1 | FLPPNRYIFHGAEVYSDEDDVLSSSSCGSNSDSGTCQSPSLEEPMED ESEIEEFYNGLE    | 703 |
| hsIRT1 | DEPDVPERAGGAGFGTDGDDQEAINEAISVKQEV TDMNYP SNKS                  | 747 |

